

**Table 5.26-- WATER QUALITY AT SELECTED PUBLIC BEACHES:
1999 AND 2000**

Island and beach	Number of samples		Enterococci density 1/	
	1999	2000	1999	2000
Hawaii:				
Hapuna Beach	38	-	0.5	(X)
Kahaluu Beach	39	2	3.7	5.1
Kealakekua Bay (off curio stand)	-	-	(X)	(X)
Hilo Bay (Mooheau Park)	-	-	(X)	(X)
Spencer Beach Park	46	35	2.6	1.3
Maui:				
Kapalua (Fleming) Beach (north)	48	31	1.0	1.0
Kihei (north)	-	-	(X)	(X)
Makena Beach	-	-	(X)	(X)
Seven Pools	3	3	0.5	0.3
Sheraton Kaanapali (shoreline)	39	-	0.8	(X)
Lanai:				
Hulopoe Bay	-	-	(X)	(X)
Molokai:				
Kaunakakai Harbor	-	-	(X)	(X)
Oahu:				
Ala Moana Park (center)	4	40	9.7	2.1
Ewa Beach Park	7	11	0.8	3.5
Haleiwa Beach	25	49	3.0	2.2
Hanauma Bay	49	51	4.2	1.5
Kailua Beach Park	50	51	1.9	3.2
Kuhio Beach	49	48	8.7	4.9
Makaha Beach	-	-	(X)	(X)
Waimea Beach	41	-	3.2	(X)
Kauai:				
Anini Park Pavilion	43	13	1.6	14.3
Kalapaki Beach (middle)	50	50	4.7	6.1
Kekaha (Oomano Point)	-	-	(X)	(X)
Poipu Beach Pavilion	43	35	1.4	1.1
West of Lydgate Park (wading pool)	50	51	1.2	2.4

X Not applicable.

1/ Geometric mean, number per 100 ml. The geometric mean standard for Enterococci density is 7 per 100 ml.

Source: Hawaii State Department of Health, Clean Water Branch, records.